

REMARKS ABOUT SCHLUMPRECHT SPACE

DENKA KUTZAROVA

ABSTRACT. We prove that Schlumprecht space S is isomorphic to $(\sum_{k=1}^{\infty} \oplus \ell_{\infty}^{n_k})_S$ for any sequence of integers (n_k) .

Th. Schlumprecht constructed the first arbitrarily distortable space S [S1]. He also proved that S is complementably minimal [S2]. A Banach space X is called minimal (a notion defined by H.P. Rosenthal) if every infinite dimensional subspace of X contains a further subspace isomorphic to X and X is called complementably minimal (defined by A. Pełczyński) if every infinite dimensional subspace of X contains a subspace which is isomorphic to X and complemented in X .

As it was remarked in [S2], the space S is either prime or fails the Schroeder-Bernstein property (see [C], [G2]). A Banach space X is called prime [LT] if every complemented infinite dimensional subspace of X is isomorphic to X . A. Pełczyński showed that the spaces ℓ_p , $1 \leq p < \infty$, and c_0 are prime, while J. Lindenstrauss proved that ℓ_{∞} is also prime (cf. e.g. [LT]). Recently W.T. Gowers and B. Maurey [GM] constructed more prime spaces for which all complemented infinite dimensional subspaces are of finite codimension. It is not known yet if S is prime. It seems hard to characterize its complemented subspaces since S is rich with them. We shall see that for every sequence $(n_k)_{k=1}^{\infty}$ of integers, the Schlumprecht sum $Y = \left(\sum_{k=1}^{\infty} \oplus \ell_{\infty}^{n_k} \right)_S$ is isomorphic to a complemented subspace of S . For some sequences, one can immediately observe that the corresponding sum is isomorphic to S . For example, if $n_k = 2^k$, we use that the standard basis of S is subsymmetric to show that Y is isomorphic to its square and then it follows by Pełczyński decomposition technique that $Y \approx S$ (see [C] for more details). We hoped in the beginning that there exists a sequence $\{n_k\}$ for which Y is not isomorphic to S and thus to show that S fails to be prime. Unfortunately, this turned out not to be true.

Recall the definition of Schlumprecht space S . Let $(e_i)_{i=1}^{\infty}$ be the standard basis of the linear space c_{00} of finitely supported sequences. For $x = \sum_{i=1}^{\infty} a_i e_i \in c_{00}$, set $\text{supp } x = \{i \in \mathbb{N} : a_i \neq 0\}$. For E, F finite subsets of $\mathbb{N}$, $E < F$ means $\max E < \min F$ or either E or F is empty. For $x = \sum_{i=1}^{\infty} a_i e_i$ and E a subset of $\mathbb{N}$, we denote by Ex the vector $Ex = \sum_{i \in E} a_i e_i$. For $t \geq 1$ let $f(t) = \log_2(t+1)$. Then $S = (S, \|\cdot\|)$ is the completion of c_{00} with respect to the norm $\|\cdot\|$ defined by the equation:

$$\|x\| = \max \left\{ \|x\|_\infty, \sup_{\substack{r \geq 2 \\ E_1 < E_2 < \dots < E_r}} \frac{1}{f(r)} \sum_{i=1}^r \|E_i x\| \right\}.$$

Clearly, (e_i) is a 1-unconditional and 1-subsymmetric basis of S .

We shall use the following results from [S2].

Statement 1. Every normalized block basis (y_i) of (e_i) dominates (e_i) , i.e.

$$\left\| \sum_{i=1}^{\infty} a_i y_i \right\| \geq \left\| \sum_{i=1}^{\infty} a_i e_i \right\|$$

for all $(a_i) \in c_{00}$.

Statement 2. (Lemma 2 [S2]) Let $\varepsilon > 0$ and $r \in \mathbb{N}$. Then there is an $n = n(\varepsilon, r) \in \mathbb{N}$ with following property: if $m \geq n$ and if $y = \frac{1}{m} \sum_{i=1}^m x_i$, where $(x_i)_{i=1}^m$ is a normalized block basis of (e_i) which is $(1 + \varepsilon/2)$ - equivalent to the unit basis of ℓ_1^m , then

$$\sup_{E_1 < E_2 < \dots < E_r} \sum_{i=1}^r \|E_i y\| \leq \|y\| + \varepsilon.$$

Statement 3. (Theorem 3 [S2]) Let $\varepsilon_k > 0, k = 1, 2, \dots$ with $\sum \varepsilon_k \leq 1$. There exists a constant $C > 1$ such that for any normalized block basis (y_k) of (e_k) with the following properties: there is a sequence $r_k \uparrow \infty$ in $\mathbb{N}$ so that for all $k \in \mathbb{N}$,

$$(a) \quad \sup_{\substack{r \leq r_{k-1} \\ E_1 < E_2 < \dots < E_r}} \sum_{i=1}^r \|E_i(y_k)\| \leq 1 + \varepsilon_k,$$

$$(b) \quad \text{card}(\text{supp } y_k) \leq \varepsilon_k \cdot f\left(\frac{r_k}{3}\right),$$

we have that (y_k) is C -equivalent to (e_k) .

The above statements easily imply that S is complementably minimal. It is known that S contains uniformly $\ell_\infty^n, n = 1, 2, \dots$. The construction of a $(1 + \delta)$ -copy of ℓ_∞^n looks like a “yardstick”. For example, let $F_1, F_2, \dots, F_n$ be disjoint subsets of $\mathbb{N}$ with $\text{card}(F_1) = p_1 = q_1$, $\text{card}(F_2) = p_2 = q_1 q_2, \dots$, $\text{card}(F_n) = p_n = q_1 q_2 \dots q_n$. We also construct $F_1, \dots, F_n$ to be “nested”, i.e. for $\chi_{F_i} = \{e_{k_j}^i\}_{j=1}^{p_i}, i = 1, 2, \dots, n$, we require $\text{supp}\{e_{k_1}^i\} < \text{supp}\{e_{k_j}^{i+1} : j = 1, 2, \dots, q_{i+1}\} < \text{supp}\{e_{k_2}^i\} < \text{supp}\{e_{k_j}^{i+1} : j = q_{i+1} + 1, \dots, 2q_{i+1}\} < \dots < \text{supp}\{e_{k_{p_i}}^i\} < \text{supp}\{e_{k_j}^{i+1} : j = (p_i - 1)q_{i+1}, \dots, p_{i+1}\}$.

If q_i are chosen big enough (depending on n and δ), $q_1 \ll \dots \ll q_n$, then the finite sequence of normalized vectors $y_i = \frac{1}{f(p_i)} \chi_{F_i}, i = 1, 2, \dots, n$, is $(1 + \delta)$ -equivalent to the unit basis of ℓ_∞^n . A proof of this fact was never written. One can find an argument in the same spirit in [ADKM]; it is more complicated, however, because mixed Tsirelson spaces are asymptotic ℓ_1 spaces.

Let now $(n_k)_{k=1}^\infty$ be an arbitrary sequence of integers. We shall construct a subspace Y of S which is isomorphic to $(\Sigma \oplus \ell_\infty^{n_k})_S$. Recall that given $(X_k, \|\cdot\|_k)$, the norm of $y = \Sigma y_k$ in the sum is defined as $\|\Sigma\|y_k\|_k e_k\|$, where $\|\cdot\|$ is the standard norm of S .

In view of Statement 3, choose a sequence of positive reals ε_k with $\Sigma \varepsilon_k \leq 1$. We shall construct successively for $k = 1, 2, \dots$ collections $(z_i^k)_{i=1}^{n_k}$ of normalized elements in S with $[(z_i^k)_{i=1}^{n_k}]$ being $(1 + \delta_k)$ -isomorphic to $\ell_\infty^{n_k}$ for sufficiently small δ_k such that $y_k = z_k / \|z_k\|$, where $z_k = \sum_{i=1}^{n_k} z_i^k$, satisfy the conditions of Statement 3. Each z_k will be an arithmetic mean of m_k “yardsticks”, m_k big enough and $m_k \ll q_1^k \ll q_2^k \ll \dots \ll q_{n_k}^k$. That is, we pick a yardstick, described above, then we take m_k successive copies of it, i.e. m_k successive vectors with the same distribution. Thus, each $z_i^k, i = 1, \dots, n_k$, is of the form from Statement 2 with $m = m_k$. Note that if $n_k = n_i$ for $k \neq i$, in order to satisfy the conditions of Statement 3, the concrete representations of $\ell_\infty^{n_k}$ and $\ell_\infty^{n_i}$ in S have different distributions.

By Statement 3, (y_k) is C -isomorphic to (e_k) . The form of y_k also assures that if for each k we take an arbitrary combination $x_k = \sum_{i=1}^{n_k} a_i^k y_i^k$, then the sequence $(x_k / \|x_k\|)$ also satisfies the hypothesis of Statement 3. Therefore, $(x_k / \|x_k\|)$ is C -isomorphic to (e_k) . Set $Y = [(y_k)_{k=1}^\infty]$. Thus, we obtained

Proposition 1. The subspace Y is $2C$ -isomorphic to $(\sum_{k=1}^\infty \oplus \ell_\infty^{n_k})_S$.

We also have

Proposition 2. The subspace Y is complemented in S .

Proof. Let E_k be intervals of integers, containing $\text{supp } y_k$ and $E_1 < E_2 < \dots$. Since the subspaces $\ell_\infty^{n_k}$ are uniformly complemented, see [LT], let Q_k be projections from $E_k S$ onto $[(y_i)_{i=1}^{n_k}]$ with $\|Q_k\| \leq 3$. It follows from Statement 1 that for every $x \in S$,

$$\left\| \sum_{k=1}^\infty \|E_k x\| e_k \right\| \leq \left\| \sum_{k=1}^\infty E_k x \right\| \leq \|x\|.$$

On the other hand, for the map $P = \sum_k Q_k E_k$ we have by Proposition 1 that

$$\|Px\| \leq 2C \left\| \sum_{k=1}^\infty \|Q_k E_k x\| e_k \right\| \leq 6C \left\| \sum_{k=1}^\infty \|E_k x\| e_k \right\|.$$

Therefore, $\|Px\| \leq 6C\|x\|$. Clearly, $P^2 = P$ and $P(S) = Y$, which ends the proof. $\square$

In order to show that Y is isomorphic to S , we shall use a variant of the refinement of Johnson’s argument, presented in [BCLT], Proposition 6.3, for a space with a subsymmetric basis. We shall work with finite dimensional decompositions instead of block-vectors.

Proposition 3. The subspace Y is isomorphic to S .

Proof. Following [BCLT], represent every integer in the form $m = 2^i(2j-1)$, $i = 0, 1, 2, \dots, j = 1, 2, \dots$. As above, we construct inductively a sequence of consecutive finite dimensional

subspaces (V_m) , i.e. $\text{supp } V_1 < \text{supp } V_2 < \dots$, where we define the support of a subspace to be the union of the supports of its elements. We construct (V_m) to be generated by an appropriate yardstick which represents in S the space $\ell_\infty^{n_j}$ and we also assure that the conditions of Statement 3 are satisfied. Note that the representation of $\ell_\infty^{n_j}$ have different distribution for different m . As in Proposition 1, the subspace $V = [(V_m)_{m=1}^\infty]$ is isomorphic to the sum $Z = (\ell_\infty^{n_1} \oplus \ell_\infty^{n_1} \oplus \ell_\infty^{n_2} \oplus \ell_\infty^{n_1} \oplus \ell_\infty^{n_3} \oplus \dots)_S$. By Proposition 2, V is complemented in S . We have that $[(V_{2m})_{m=1}^\infty]$ is also isomorphic to Z and therefore, it is also isomorphic to V . The subspace $[(V_{2k-1})_{k=1}^\infty]$ is isomorphic to $(\sum_{k=1}^\infty \oplus \ell_\infty^{n_k})_S$ and hence it is isomorphic to Y . Since V is complemented, $S \approx V \oplus W$, whence

$$Y \oplus S \approx Y \oplus V \oplus W \approx V \oplus W \approx S.$$

Since S is complementably minimal, $Y \approx S \oplus X$. Evidently $S \oplus S \approx S$ and therefore,

$$Y \approx S \oplus X \approx S \oplus S \oplus X \approx S \oplus Y \approx S.$$

Thus, $Y \approx S$, which implies $S \approx (\Sigma \oplus \ell_\infty^{n_k})_S$. $\square$

A Banach space with an unconditional basis is said to have a unique unconditional basis if any two normalized unconditional bases are equivalent after a permutation, see [BCLT] and for recent results see [CK]. It was remarked in [CK] that the example of Gowers in [G1] has a unique unconditional basis. Proposition 3 implies that S has no unique unconditional basis. This also immediately follows if one uses that S is complementably minimal and combines Proposition 6.3 and Proposition 6.4 [BCLT]. They provide the existence of a normalized block basis with constant coefficients (u_i) which spans a subspace isomorphic to S and no permutation of (u_i) is equivalent to (e_i) . In fact, one can construct such a block-basis (u_i) directly without Proposition 6.4 which uses an adaptation of Zippin's argument. Indeed, if we consider consecutive blocks with constant coefficients whose length increases rapidly enough, it is easy to show that Statement 3 and Proposition 6.3 [BCLT] imply the following.

Proposition 4. Schlumprecht space S is isomorphic to $(\sum_k \oplus \ell_1^{n_k})_S$ for any sequence of integers (n_k) .

P. Casazza pointed out the following immediate consequence of our observations (in fact, not using their full generality). Recall that the same property for the classical spaces ℓ_p for $p > 2$, $1 < p < 2$ and $p = 1$ was shown in [R], [BDGJN] and [B] respectively.

Corollary 5. Schlumprecht space S (resp. its dual S^*) has a subspace isomorphic to the whole space and not complemented in S (resp. S^*).

One possible way to see it is to use the fact that one can embed ℓ_1^n in ℓ_∞^m with a bad constant of complementation.

Remark. Let Z be a complemented subspace of S which has a subsymmetric basis. Since Z is isomorphic to its square, then by the decomposition technique we have that Z is isomorphic to S .

Question. Does S have a unique subsymmetric basis?

REFERENCES

- [ADKM] S.A. Argyros, I. Deliyanni, D.N. Kutzarova and A. Manoussakis, Modified mixed Tsirelson spaces, preprint.
- [BDGJN] B. Bennett, L.E. Dor, V. Goodman, W.B. Johnson and C.M. Newman, On uncomplemented subspaces of L_p , $1 < p < 2$, Israel J. Math. 26 (1977), 178-187.
- [B] J. Bourgain, A counterexample to a complementation problem, Compositio Math. 43 (1981), 133-144.
- [BCLT] J. Bourgain, P.G. Casazza, J. Lindenstrauss and L. Tzafriri, Banach spaces with a unique unconditional basis, up to a permutation, Memoirs Amer. Math. Soc. 322, 1985.
- [C] P.G. Casazza, The Schroeder-Bernstein property for Banach spaces, Contemp. Math., 85 (1989), 61-77.
- [CK] P.G. Casazza and N.J. Kalton, Uniqueness of unconditional bases in Banach spaces, Israel J. Math., to appear.
- [G1] W.T. Gowers, A solution to Banach's hyperplane problem, Bull. London Math. Soc. 26 (1994), 523-530.
- [G2] W.T. Gowers, A solution to the Schroeder-Bernstein problem for Banach spaces, Bull. London Math. Soc. 28 (1996), 297-304.
- [GM] W.T. Gowers and B. Maurey, Banach spaces with small spaces of operators, Math. Annal. 307 (1997), 543-568.
- [LT] J. Lindenstrauss and L. Tzafriri, Classical Banach spaces I, Sequence Spaces, Springer, 1977.
- [R] H.P. Rosenthal, on the subspaces of L^p , $p > 2$ spanned by sequences of independent random variables, Israel J. Math. 8 (1970), 273-303.
- [S1] Th. Schlumprecht, An arbitrarily distortable Banach space, Israel J. Math. 76 (1991), 81-95.
- [S2] Th. Schlumprecht, A complementably minimal Banach space not containing c_0 or ℓ_p , preprint.

INSTITUTE OF MATHEMATICS, BULGARIAN ACADEMY OF SCIENCES, 1113 SOFIA, BULGARIA
Current address: Miami University, Oxford, Ohio, USA, e-mail address: kutzard@muohio.edu